

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/122644/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

El-Hiti, Gamal A., Abdel-Wahab, Bakr F., Yousif, Emad, Hayal Alotaibi, Mohammad, Hegazy, Amany S. and Kariuki, Benson M. ORCID: <https://orcid.org/0000-0002-8658-3897> 2019. 5-[5-(4-Chlorophenyl)isoxazol-3-yl]-N-phenyl-1,3,4-oxadiazol-2-amine. IUCrData 4 (1) , x190136. 10.1107/S2414314619001366 file

Publishers page: <http://dx.doi.org/10.1107/S2414314619001366>
<<http://dx.doi.org/10.1107/S2414314619001366>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.





IUCrData

ISSN 2414-3146

5-[5-(4-Chlorophenyl)isoxazol-3-yl]-N-phenyl-1,3,4-oxadiazol-2-amine

Gamal A. El-Hiti,^{a*} Bakr F. Abdel-Wahab,^{b,c} Emad Yousif,^d Mohammad Hayal Alotaibi,^e Amany S. Hegazy^f and Benson M. Kariuki^{f‡}

^aDepartment of Optometry, College of Applied Medical Sciences, King Saud, University, PO Box 10219, Riyadh 11433, Saudi Arabia, ^bDepartment of Chemistry, College of Science and Humanities, Shaqra University, Duwadimi, Saudi Arabia, ^cApplied Organic Chemistry Department, National Research Centre, Dokki, Giza 12622, Egypt, ^dDepartment of Chemistry, College of Science, Al-Nahrain University, Baghdad 64021, Iraq, ^eNational Center for Petrochemicals Technology, King Abdulaziz City for Science and Technology, PO Box 6086, Riyadh 11442, Saudi Arabia, and ^fSchool of Chemistry, Cardiff University, Main Building, Park Place, Cardiff CF10 3AT, UK. *Correspondence e-mail: gelhiti@ksu.edu.sa

Received 23 January 2019

Accepted 24 January 2019

Edited by W. T. A. Harrison, University of Aberdeen, Scotland

‡ Additional corresponding author, e-mail: kariukib@cardiff.ac.uk.


Keywords: crystal structure; oxadiazole; isoxazole.

CCDC reference: 1893181


Structural data: full structural data are available from iucrdata.iucr.org

The title compound, C₁₇H₁₁ClN₄O₂, consists of chlorophenyl (A), isoxazolyl (B), oxadiazolyl (C) and phenyl (D) rings with twist angles between the planes through neighbouring rings A/B, B/C and C/D of 14.9 (1), 1.8 (1) and 8.9 (1)° respectively. In the crystal, adjacent molecules related by inversion symmetry form columns parallel to the [010] direction through π - π stacking [shortest centroid-centroid separation = 3.6203 (10) Å]; these are cross-linked by N—H...N interactions in the [001] direction.

3D view



Chemical scheme




Structure description

Substituted 1,3,4-oxadiazol-2-amines can act as anticancer and antiproliferative agents (Ahsan *et al.*, 2014; Gonda *et al.*, 2017; Kumar *et al.*, 2009, 2011). As part of our studies in this area, we now describe the synthesis and structure of the title compound.

The molecule consists of chlorophenyl (A), isoxazolyl (B), oxadiazolyl (C) and phenyl (D) rings (Fig. 1). The molecule is close to planar, as indicated by the twist angles between the planes through neighbouring rings A/B, B/C and C/D, namely 14.9 (1), 1.8 (1) and 8.9 (1)° respectively. A short intramolecular C—H...N contact is observed (Table 1). In the crystal, adjacent molecules related by inversion symmetry form columns parallel to the [010] direction through π - π stacking [shortest centroid-centroid separation = 3.6203 (10) Å]. The columns of molecules are linked by N—H...N interactions in the [001] direction (Fig. 2).

Table 1
Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$C17-H17\cdots N3$	0.93	2.33	2.955 (2)	125
$N4-H4\cdots N2^i$	0.86	2.42	3.176 (2)	146


Symmetry code: (i) $x, -y + \frac{3}{2}, z - \frac{1}{2}$.**Figure 1**
The molecular structure of the title compound showing 50% probability ellipsoids.

Synthesis and crystallization

2-(3-(4-Chlorophenyl)isoxazole-5-carbonyl)-*N*-phenylhydrazinecarbothioamide was treated with anhydrous sodium acetate in ethanol under reflux for 5 h. The resulting solid was collected by filtration, washed with ethanol, dried and recrystallized from dimethylformamide solution to give colourless plates, m.p. 277–278°C.

Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2.

**Figure 2**
A segment of the crystal structure showing $N-H\cdots N$ hydrogen bonds as dotted lines.**Table 2**
Experimental details.

Crystal data	
Chemical formula	$C_{17}H_{11}ClN_4O_2$
M_r	338.75
Crystal system, space group	Monoclinic, $P2_1/c$
Temperature (K)	296
a, b, c (Å)	17.8925 (15), 7.4168 (6), 11.5006 (9)
β (°)	102.061 (8)
V (Å ³)	1492.5 (2)
Z	4
Radiation type	Mo $K\alpha$
μ (mm ⁻¹)	0.27
Crystal size (mm)	0.32 × 0.17 × 0.04
Data collection	
Diffractometer	Rigaku Oxford Diffraction SuperNova, Dual, Cu at zero, Atlas Gaussian (<i>CrysAlis PRO</i> ; Rigaku OD, 2015)
Absorption correction	
T_{min}, T_{max}	0.768, 1.000
No. of measured, independent and observed [$I > 2\sigma(I)$] reflections	13114, 3720, 2622
R_{int}	0.027
$(\sin \theta/\lambda)_{max}$ (Å ⁻¹)	0.699
Refinement	
$R[F^2 > 2\sigma(F^2)], wR(F^2), S$	0.045, 0.112, 1.06
No. of reflections	3720
No. of parameters	218
H-atom treatment	H-atom parameters constrained
$\Delta\rho_{max}, \Delta\rho_{min}$ (e Å ⁻³)	0.25, -0.23

Computer programs: *CrysAlis PRO* (Rigaku OD, 2015), *SHELXS* (Sheldrick, 2008), *SHELXL2018* (Sheldrick, 2015), *ORTEP-3 for Windows* and *WinGX* (Farrugia, 2012) and *CHEMDRAW Ultra* (Cambridge Soft, 2001).

Funding information

MHA thanks King Abdulaziz City for Science and Technology (KACST), Saudi Arabia for financial support (award No. 020-0180).

References

- Ahsan, M. J., Sharma, J., Singh, M., Jadav, S. S. & Yasmin, S. (2014). *Biomed. Res. Int.* **2014**, 814984.
- Cambridge Soft (2001). *CHEMDRAW Ultra*. Cambridge Soft Corporation, Cambridge, Massachusetts, USA.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Gonda, T., Bérđi, P., Zupkó, I., Fülöp, F. & Szakonyi, Z. (2017). *Int. J. Mol. Sci.* **19**, 81.
- Kumar, D., Patel, G., Chavers, A. K., Chang, K. H. & Shah, K. (2011). *Eur. J. Med. Chem.* **46**, 3085–3092.
- Kumar, D., Patel, G., Johnson, E. O. & Shah, K. (2009). *Bioorg. Med. Chem. Lett.* **19**, 2739–2741.
- Rigaku OD (2015). *CrysAlis PRO*. Rigaku Oxford Diffraction, Yarnton, England.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.

full crystallographic data

IUCrData (2019). 4, x190136 [https://doi.org/10.1107/S2414314619001366]

5-[5-(4-Chlorophenyl)isoxazol-3-yl]-*N*-phenyl-1,3,4-oxadiazol-2-amine

Gamal A. El-Hiti, Bakr F. Abdel-Wahab, Emad Yousif, Mohammad Hayal Alotaibi, Amany S. Hegazy and Benson M. Kariuki

5-[5-(4-Chlorophenyl)isoxazol-3-yl]-*N*-phenyl-1,3,4-oxadiazol-2-amine*Crystal data*

$C_{17}H_{11}ClN_4O_2$

$M_r = 338.75$

Monoclinic, $P2_1/c$

$a = 17.8925$ (15) Å

$b = 7.4168$ (6) Å

$c = 11.5006$ (9) Å

$\beta = 102.061$ (8)°

$V = 1492.5$ (2) Å³

$Z = 4$

$F(000) = 696$

$D_x = 1.508$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3445 reflections

$\theta = 3.3$ – 26.7°

$\mu = 0.27$ mm⁻¹

$T = 296$ K

Plate, colourless

$0.32 \times 0.17 \times 0.04$ mm

Data collection

Rigaku Oxford Diffraction SuperNova, Dual,
Cu at zero, Atlas
diffractometer

ω scans

Absorption correction: gaussian
(CrysAlisPro; Rigaku OD, 2015)

$T_{\min} = 0.768$, $T_{\max} = 1.000$

13114 measured reflections

3720 independent reflections

2622 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.027$

$\theta_{\max} = 29.8^\circ$, $\theta_{\min} = 3.0^\circ$

$h = -24 \rightarrow 23$

$k = -10 \rightarrow 10$

$l = -13 \rightarrow 16$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.045$

$wR(F^2) = 0.112$

$S = 1.06$

3720 reflections

218 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.039P)^2 + 0.4804P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.25$ e Å⁻³

$\Delta\rho_{\min} = -0.23$ e Å⁻³

Extinction correction: SHELXL-2018/1
(Sheldrick 2018),

$F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.0044 (7)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.27509 (9)	0.6433 (2)	0.57408 (14)	0.0354 (4)
C2	0.27554 (10)	0.5711 (2)	0.68585 (14)	0.0380 (4)
H2	0.321696	0.540461	0.735750	0.046*
C3	0.20791 (11)	0.5447 (2)	0.72307 (15)	0.0406 (4)
H3	0.208175	0.496146	0.797718	0.049*
C4	0.13988 (10)	0.5911 (2)	0.64829 (16)	0.0419 (4)
C5	0.13775 (10)	0.6597 (3)	0.53621 (16)	0.0443 (4)
H5	0.091345	0.687890	0.486141	0.053*
C6	0.20539 (10)	0.6856 (2)	0.49975 (15)	0.0410 (4)
H6	0.204587	0.732140	0.424398	0.049*
C7	0.34717 (10)	0.6763 (2)	0.53732 (14)	0.0368 (4)
C8	0.42112 (10)	0.6850 (2)	0.59279 (15)	0.0411 (4)
H8	0.441861	0.666766	0.673050	0.049*
C9	0.45996 (10)	0.7280 (2)	0.50222 (15)	0.0395 (4)
C10	0.54121 (10)	0.7559 (2)	0.50955 (14)	0.0386 (4)
C11	0.64134 (9)	0.8153 (2)	0.44570 (14)	0.0363 (4)
C12	0.76067 (9)	0.8721 (2)	0.37782 (14)	0.0348 (4)
C13	0.78904 (10)	0.9338 (2)	0.28169 (15)	0.0416 (4)
H13	0.755572	0.967507	0.211870	0.050*
C14	0.86646 (11)	0.9450 (3)	0.28962 (17)	0.0480 (5)
H14	0.885249	0.984768	0.224560	0.058*
C15	0.91682 (11)	0.8977 (3)	0.39347 (17)	0.0494 (5)
H15	0.969268	0.906781	0.398840	0.059*
C16	0.88857 (10)	0.8372 (3)	0.48854 (17)	0.0473 (5)
H16	0.922266	0.805371	0.558565	0.057*
C17	0.81049 (10)	0.8229 (2)	0.48169 (15)	0.0392 (4)
H17	0.791856	0.780592	0.546287	0.047*
N1	0.41386 (9)	0.7429 (3)	0.39846 (14)	0.0606 (5)
N2	0.59647 (9)	0.7436 (2)	0.59928 (13)	0.0490 (4)
N3	0.66384 (8)	0.7833 (2)	0.55885 (13)	0.0487 (4)
N4	0.68044 (8)	0.8585 (2)	0.36244 (12)	0.0405 (4)
H4	0.654043	0.880249	0.292363	0.049*
O1	0.34130 (7)	0.7092 (2)	0.42024 (11)	0.0600 (4)
O2	0.56429 (6)	0.80070 (17)	0.40728 (10)	0.0409 (3)
Cl1	0.05523 (3)	0.56575 (9)	0.69766 (5)	0.0682 (2)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0365 (9)	0.0360 (9)	0.0339 (8)	−0.0021 (7)	0.0078 (7)	−0.0019 (7)
C2	0.0399 (9)	0.0403 (9)	0.0324 (8)	−0.0005 (7)	0.0046 (7)	−0.0025 (7)
C3	0.0490 (11)	0.0427 (10)	0.0317 (8)	−0.0024 (8)	0.0119 (7)	−0.0009 (8)
C4	0.0389 (10)	0.0444 (10)	0.0465 (10)	−0.0004 (8)	0.0178 (8)	−0.0048 (8)
C5	0.0372 (10)	0.0494 (11)	0.0455 (10)	0.0037 (8)	0.0069 (8)	0.0012 (9)
C6	0.0422 (10)	0.0467 (10)	0.0344 (9)	0.0018 (8)	0.0088 (7)	0.0054 (8)
C7	0.0375 (9)	0.0409 (9)	0.0315 (8)	−0.0010 (7)	0.0061 (7)	0.0015 (7)
C8	0.0373 (9)	0.0538 (11)	0.0316 (8)	−0.0017 (8)	0.0062 (7)	−0.0005 (8)
C9	0.0346 (9)	0.0462 (10)	0.0373 (9)	−0.0011 (8)	0.0068 (7)	0.0013 (8)
C10	0.0348 (9)	0.0480 (10)	0.0338 (8)	−0.0012 (8)	0.0087 (7)	0.0001 (8)
C11	0.0288 (8)	0.0430 (9)	0.0362 (9)	0.0006 (7)	0.0046 (7)	−0.0043 (7)
C12	0.0326 (9)	0.0356 (9)	0.0357 (8)	−0.0002 (7)	0.0063 (7)	−0.0050 (7)
C13	0.0367 (9)	0.0515 (11)	0.0355 (9)	−0.0013 (8)	0.0049 (7)	0.0017 (8)
C14	0.0435 (11)	0.0560 (12)	0.0471 (11)	−0.0051 (9)	0.0154 (8)	0.0002 (9)
C15	0.0327 (9)	0.0614 (12)	0.0544 (11)	−0.0017 (9)	0.0101 (8)	−0.0043 (10)
C16	0.0372 (10)	0.0564 (12)	0.0443 (10)	0.0034 (9)	−0.0003 (8)	−0.0018 (9)
C17	0.0380 (9)	0.0444 (10)	0.0344 (9)	0.0005 (8)	0.0059 (7)	−0.0010 (8)
N1	0.0337 (9)	0.1058 (15)	0.0422 (9)	−0.0077 (9)	0.0076 (7)	0.0204 (9)
N2	0.0364 (8)	0.0768 (12)	0.0341 (8)	−0.0046 (8)	0.0084 (6)	0.0010 (8)
N3	0.0349 (8)	0.0784 (12)	0.0323 (8)	−0.0047 (8)	0.0059 (6)	−0.0001 (8)
N4	0.0304 (7)	0.0583 (9)	0.0316 (7)	0.0004 (7)	0.0035 (6)	0.0050 (7)
O1	0.0337 (7)	0.1065 (12)	0.0386 (7)	−0.0082 (7)	0.0048 (5)	0.0204 (7)
O2	0.0308 (6)	0.0564 (8)	0.0345 (6)	−0.0015 (5)	0.0050 (5)	0.0041 (6)
Cl1	0.0494 (3)	0.0880 (4)	0.0766 (4)	0.0034 (3)	0.0345 (3)	0.0077 (3)

Geometric parameters (\AA , $^\circ$)

C1—C2	1.391 (2)	C10—O2	1.3665 (19)
C1—C6	1.393 (2)	C11—N3	1.301 (2)
C1—C7	1.459 (2)	C11—N4	1.338 (2)
C2—C3	1.379 (2)	C11—O2	1.3611 (19)
C2—H2	0.9300	C12—C17	1.382 (2)
C3—C4	1.379 (3)	C12—C13	1.387 (2)
C3—H3	0.9300	C12—N4	1.413 (2)
C4—C5	1.379 (3)	C13—C14	1.372 (3)
C4—Cl1	1.7351 (18)	C13—H13	0.9300
C5—C6	1.375 (2)	C14—C15	1.383 (3)
C5—H5	0.9300	C14—H14	0.9300
C6—H6	0.9300	C15—C16	1.372 (3)
C7—C8	1.345 (2)	C15—H15	0.9300
C7—O1	1.351 (2)	C16—C17	1.387 (2)
C8—C9	1.404 (2)	C16—H16	0.9300
C8—H8	0.9300	C17—H17	0.9300
C9—N1	1.306 (2)	N1—O1	1.3952 (19)
C9—C10	1.453 (2)	N2—N3	1.411 (2)

C10—N2	1.275 (2)	N4—H4	0.8600
C2—C1—C6	119.05 (16)	N3—C11—N4	131.36 (16)
C2—C1—C7	119.83 (15)	N3—C11—O2	113.02 (15)
C6—C1—C7	121.11 (15)	N4—C11—O2	115.62 (14)
C3—C2—C1	120.36 (16)	C17—C12—C13	119.92 (16)
C3—C2—H2	119.8	C17—C12—N4	122.98 (15)
C1—C2—H2	119.8	C13—C12—N4	117.08 (15)
C4—C3—C2	119.23 (16)	C14—C13—C12	119.99 (17)
C4—C3—H3	120.4	C14—C13—H13	120.0
C2—C3—H3	120.4	C12—C13—H13	120.0
C5—C4—C3	121.56 (16)	C13—C14—C15	120.58 (18)
C5—C4—C11	119.36 (14)	C13—C14—H14	119.7
C3—C4—C11	119.07 (14)	C15—C14—H14	119.7
C6—C5—C4	118.86 (17)	C16—C15—C14	119.29 (17)
C6—C5—H5	120.6	C16—C15—H15	120.4
C4—C5—H5	120.6	C14—C15—H15	120.4
C5—C6—C1	120.92 (16)	C15—C16—C17	120.94 (17)
C5—C6—H6	119.5	C15—C16—H16	119.5
C1—C6—H6	119.5	C17—C16—H16	119.5
C8—C7—O1	109.14 (15)	C12—C17—C16	119.28 (16)
C8—C7—C1	135.40 (16)	C12—C17—H17	120.4
O1—C7—C1	115.45 (14)	C16—C17—H17	120.4
C7—C8—C9	104.44 (15)	C9—N1—O1	104.75 (14)
C7—C8—H8	127.8	C10—N2—N3	106.85 (14)
C9—C8—H8	127.8	C11—N3—N2	105.10 (13)
N1—C9—C8	112.40 (16)	C11—N4—C12	126.68 (14)
N1—C9—C10	118.31 (15)	C11—N4—H4	116.7
C8—C9—C10	129.29 (16)	C12—N4—H4	116.7
N2—C10—O2	113.04 (15)	C7—O1—N1	109.26 (13)
N2—C10—C9	129.36 (16)	C11—O2—C10	101.99 (13)
O2—C10—C9	117.60 (14)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C17—H17 \cdots N3	0.93	2.33	2.955 (2)	125
N4—H4 \cdots N2 ⁱ	0.86	2.42	3.176 (2)	146

Symmetry code: (i) $x, -y+3/2, z-1/2$.